FROM POINCARÉ INEQUALITIES TO NONLINEAR MATRIX CONCENTRATION

DE HUANG AND JOEL A. TROPP

ABSTRACT. This paper deduces exponential matrix concentration from a Poincaré inequality via a short, conceptual argument. Among other examples, this theory applies to matrix-valued functions of a uniformly log-concave random vector. The proof relies on the subadditivity of Poincaré inequalities and a chain rule inequality for the trace of the matrix Dirichlet form. It also uses a symmetrization technique to avoid difficulties associated with a direct extension of the classic scalar argument.

1. Introduction

Matrix concentration inequalities describe the probability that a random matrix is close to its expected value, with deviations measured by the ℓ_2 operator norm. These results have had a profound impact on a wide range of areas in computational mathematics and statistics. See the monograph [Tro15] for an introduction to the subject and its applications.

As did the field of scalar concentration, matrix concentration theory began with simple models, such as sums of independent random matrices [LP86, Rud99, AW02, Tro12] and matrix-valued martingale sequences [PX97, Olio9, Tro11]. In recent years, researchers have sought to develop results that hold for a wider class of random matrix models. Some initial successful efforts were based on exchangeable pairs techniques [MJC⁺14, PMT16], but these methods do not address all cases of interest.

Researchers have also tried to extend scalar concentration techniques based on functional inequalities. An early attempt, by Chen & Tropp [CT14], demonstrates that (traces of) matrix variance and entropy quantities are subadditive, which leads to some Poincaré and modified log-Sobolev inequalities. A number of authors, including [CH16, CHT17, CH19], have pursued this line of work. Unfortunately, these approaches have not been sufficient to reproduce all the results that have been established in the simpler models.

Very recently, Aoun et al. [ABY19] have shown that a matrix form of the Poincaré inequality implies subexponential concentration of a random matrix with respect to the ℓ_2 operator norm. We believe that this is the first instance where a matrix functional inequality leads unconditionally to a matrix concentration result (with respect to the operator norm). Nevertheless, it remains an open question to deduce a full spectrum of matrix concentration results from matrix functional inequalities.

In this paper, we improve on [ABY19] by demonstrating that the ordinary scalar Poincaré inequality also leads to subexponential concentration with respect to the operator norm. Our argument has some elements in common with the work in [ABY19], but we have found a route to avoid most of the technical difficulty of their approach.

The basic idea is to bound the trace of an odd function (for example, the hyperbolic sine) of the random matrix using a symmetrization argument. The variance appears, and the Poincaré inequality yields a bound on the variance in terms of the Dirichlet form. Last, we apply a new matrix chain rule

1

Date: 26 April 2020. Revised 11 June 2020 and 10 October 2020.

²⁰¹⁰ Mathematics Subject Classification. Primary: 60B20, 46N30. Secondary: 60J25, 46L53.

Key words and phrases. Concentration inequality; functional inequality; Markov process; matrix concentration; Poincaré inequality; semigroup.

inequality for the Dirichlet form to obtain a moment comparison. After this paper was written, we learned that Bobkov & Ledoux proposed a similar argument in the scalar case [BL97, Sec. 4].

As in the scalar setting, Poincaré inequalities may not capture the strongest concentration properties that are possible. In a companion paper [HT20], we demonstrate that *local* Poincaré inequalities lead to the optimal subgaussian concentration results. The analysis in [HT20] captures most of the previous results on matrix concentration, but it involves more technical machinery. We also refer the reader to work of Junge & Zeng [JZ15] that contains similar results in the fully noncommutative setting.

An expert reader may still wonder about the role of (modified) log-Sobolev inequalities in establishing matrix concentration inequalities. At the time of writing, it is not clear how to obtain a matrix analog of the log-Sobolev inequality that would imply matrix concentration results in the same spirit as the ones in this paper or the related works [JZ15, ABY19, HT20].

2. Main result

This section summarizes our notation and the setting for our problem. It highlights our primary result on matrix concentration, and it gives a number of examples. In the next section, we comment on the relationship with previous work.

2.1. **Notation.** Let \mathbb{H}_d be the real linear space of $d \times d$ self-adjoint complex matrices, equipped with the ℓ_2 operator norm $\|\cdot\|$. We work with both the ordinary trace, tr, and the normalized trace, $\bar{\mathrm{tr}} := d^{-1}$ tr on the space \mathbb{H}_d . Matrices, and occasionally vectors, are written in boldface italic. In particular, \boldsymbol{f} and \boldsymbol{g} refer to functions taking values in \mathbb{H}_d . The cone \mathbb{H}_d^+ contains the positive semidefinite matrices, and the symbol \leq refers to the semidefinite order.

Given a function $\varphi : \mathbb{R} \to \mathbb{R}$ taking real values, we extend it to a function $\varphi : \mathbb{H}_d \to \mathbb{H}_d$ on self-adjoint matrices by means of the spectral resolution:

$$\textbf{\textit{A}} = \sum\nolimits_{\lambda \in \operatorname{spec}(\textbf{\textit{A}})} \lambda \, \textbf{\textit{P}}_{\lambda} \in \mathbb{H}_{d} \quad \text{implies} \quad \varphi(\textbf{\textit{A}}) = \sum\nolimits_{\lambda \in \operatorname{spec}(\textbf{\textit{A}})} \varphi(\lambda) \, \textbf{\textit{P}}_{\lambda} \in \mathbb{H}_{d}.$$

Whenever we apply a scalar function, such as a power or a hyperbolic function, to a matrix, we are referring to the standard matrix function. Nonlinear functions bind before the trace.

We use familiar notation from probability. The operator \mathbb{E} returns the expectation, and $\mathbb{P}\left\{\cdot\right\}$ is the probability of an event. The symbol \sim means "has the distribution." Nonlinear functions bind before the expectation.

2.2. **Random matrices.** Let Ω be a Polish space, equipped with a probability measure μ , and write \mathbb{E}_{μ} for the integral with respect to the measure μ . Consider a μ -integrable matrix-valued function $f: \Omega \to \mathbb{H}_d$ on the state space Ω . By drawing a random variable $Z \sim \mu$, we can construct a random matrix f(Z). Our goal is to understand the concentration of f(Z) around its mean $\mathbb{E}_{\mu} f$.

Example 2.1 (Gaussians). Consider the Gaussian space (\mathbb{R}^n, γ_n) of n-dimensional real vectors equipped with the standard normal measure γ_n . Suppose we are interested in a matrix-valued function f(X) of a standard normal random vector $X \sim \gamma_n$. A familiar example [Tro15, Chap. 5] is the matrix Gaussian series

$$f(X) = \sum_{i=1}^{n} X_i A_i$$
 where $X \sim \gamma_n$ and $A_1, \dots, A_n \in \mathbb{H}_d$ are fixed. (2.1)

We will use the Gaussian case as a running example to illustrate the concepts that we introduce.

2.3. **Markov processes.** Suppose that we can identify an ergodic, reversible, time-homogeneous Markov process $(Z_t:t\geq 0)\subset \Omega$ with initial value Z_0 and stationary measure μ . This induces a matrix-valued Markov process $(f(Z_t):t\geq 0)\subset \mathbb{H}_d$.

By ergodicity, for any point $z \in \Omega$, we have the limit $\mathbb{E}[f(Z_t) | Z_0 = z] \to \mathbb{E}_{\mu} f$ as $t \to \infty$. The results in this paper build on the intuition that a random matrix f(Z) with $Z \sim \mu$ concentrates sharply about its mean when the matrix-valued Markov process tends quickly to equilibrium.

Example 2.2 (Gaussians). We can construct a reversible Markov process $(X_t : t \ge 0) \subset \mathbb{R}^n$, called the *Ornstein–Uhlenbeck (OU) process*, by means of the stochastic differential equation

$$dX_t = -X_t dt + \sqrt{2} dB_t$$
 with initial value $X_0 \in \mathbb{R}^n$,

where $(B_t : t \ge 0) \subset \mathbb{R}^n$ is random vector whose coordinates are independent Brownian motions. The stationary measure of the OU process is the standard normal distribution γ_n .

2.4. **Derivatives and energy.** To understand how quickly a matrix Markov process $f(Z_t)$ converges to stationarity, we introduce notions of the "squared derivative" and the "energy" of the function f. Inspired by [ABY19], we define the matrix *carré du champ operator* by the formula

$$\Gamma(\boldsymbol{f})(z) := \lim_{t \downarrow 0} \frac{1}{2t} \mathbb{E}\left[\left(\boldsymbol{f}(Z_t) - \boldsymbol{f}(Z_0)\right)^2 \middle| Z_0 = z\right] \in \mathbb{H}_d^+ \text{ for } z \in \Omega.$$
 (2.2)

In many instances, the carré du champ $\Gamma(f)$ has a natural interpretation as a squared derivative of f. The expectation of the carré du champ is called the *matrix Dirichlet form*:

$$\mathscr{E}(\boldsymbol{f}) := \lim_{t \downarrow 0} \frac{1}{2t} \mathbb{E}_{Z \sim \mu} \left[\left(\boldsymbol{f}(Z_t) - \boldsymbol{f}(Z_0) \right)^2 \middle| Z_0 = Z \right] \in \mathbb{H}_d^+. \tag{2.3}$$

The Dirichlet form $\mathscr{C}(f)$ reflects the total energy of the function f.

In a general setting, it requires some care to make sense of the definitions (2.2) and (2.3). Without further comment, we restrict our attention to a "nice" class of functions where the limit in (2.2) exists pointwise and in $L_1(\mu)$ and where calculus operations are justified. By approximation, our main results on concentration hold for a wider class of functions.

Example 2.3 (Gaussians). According to [ABY19, Prop. 5.5], the matrix carré du champ operator and matrix Dirichlet form of the OU process are determined by

$$\Gamma(f)(x) = \sum_{i=1}^n (\partial_i f(x))^2$$
 for $x \in \mathbb{R}^n$ and $\mathscr{E}(f) = \sum_{i=1}^n \mathbb{E}_{\gamma_n} (\partial_i f)^2$.

The interpretations as a squared derivative and an energy are evident, and it is easy to check when the carré du champ is defined.

The matrix Gaussian series (2.1) provides an illustration:

$$\Gamma(\mathbf{f}) = \mathscr{E}(\mathbf{f}) = \sum_{i=1}^{n} A_i^2.$$

This quantity is familiar from work on matrix concentration for Gaussian series [Tro15, Chap. 4].

2.5. **Trace Poincaré inequalities.** The *matrix variance* of a function $f: \Omega \to \mathbb{H}_d$ with respect to the distribution μ is defined as

$$\operatorname{Var}_{u}[f] := \mathbb{E}_{u}\left[(f - \mathbb{E}_{u} f)^{2} \right] = \mathbb{E}_{u}[f^{2}] - (\mathbb{E}_{u} f)^{2} \in \mathbb{H}_{d}^{+}. \tag{2.4}$$

As in the scalar case, the variance reflects fluctuations of the random matrix f(Z) about its mean, where the random variable $Z \sim \mu$.

We say that the Markov process satisfies a *trace Poincaré inequality* with constant $\alpha > 0$ if

$$\operatorname{tr} \operatorname{Var}_{\mu}[f] \leq \alpha \cdot \operatorname{tr} \mathscr{C}(f) \quad \text{for all } f: \Omega \to \mathbb{H}_d.$$
 (2.5)

In other words, the trace variance of f(Z) is controlled by the energy in the function f. The inequality (2.5) provides a way to quantify the ergodicity of the Markov process.

As it happens, the trace Poincaré inequality is equivalent to an ordinary Poincaré inequality. We are grateful to Ramon Van Handel for this observation. The same result has recently appeared in the independent work of Garg et al. [GKS20].

Proposition 2.4 (Equivalence of Poincaré inequalities). *Consider a Markov process* $(Z_t : t \ge 0) \subset \Omega$ *with stationary measure* μ . *The following are equivalent:*

- (1) **Scalar Poincaré.** For all $f: \Omega \to \mathbb{R}$, it holds that $\operatorname{Var}_{\mu}[f] \leq \alpha \cdot \mathscr{C}(f)$.
- (2) Trace Poincaré. For all $d \in \mathbb{N}$ and all $f : \Omega \to \mathbb{H}_d$, it holds that $\operatorname{tr} \operatorname{Var}_{\mu}[f] \leq \alpha \cdot \operatorname{tr} \mathscr{E}(f)$.

The Poincaré constant $\alpha \geq 0$ is the same for both cases.

Proof. It is evident that the validity of the trace Poincaré inequality for all $d \in \mathbb{N}$ implies the scalar Poincaré inequality. For the reverse implication, it suffices to consider a real matrix-valued function $f: \Omega \to \mathbb{H}_d(\mathbb{R})$ with zero mean. For vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^d$, define the scalar function $g(z) = \langle \mathbf{u}, \mathbf{f}(z) \mathbf{v} \rangle \in \mathbb{R}$. Apply the scalar Poincaré inequality to g and invoke the definition (2.3) of the Dirichlet form. Thus,

$$\mathbb{E}_{\mu} \left\langle \boldsymbol{u}, \, \boldsymbol{f}(z) \, \boldsymbol{v} \right\rangle^{2} \leq \alpha \cdot \lim_{t \downarrow 0} \frac{1}{2t} \, \mathbb{E}_{Z \sim \mu} \left[\left\langle \boldsymbol{u}, \, \left(\boldsymbol{f}(Z_{t}) - \boldsymbol{f}(Z_{0}) \right) \boldsymbol{v} \right\rangle^{2} \, \Big| \, Z_{0} = Z \right].$$

Instate this inequality with $\mathbf{v} = \mathbf{e}_i$ for each $i = 1, \dots, d$ and sum over i to arrive at

$$\langle u, \operatorname{Var}_{\mu}[f] u \rangle \leq \langle u, \mathscr{C}(f) u \rangle.$$

Average over $\boldsymbol{u} \sim \text{uniform}\{\pm 1\}^d$ to reach the trace Poincaré inequality. To extend this argument to complex matrices, apply the same approach to the real and imaginary parts of the inner product. \Box

The main result of this paper is that concentration properties of the random matrix f(Z) follow from the trace Poincaré inequality (2.5) or, equivalently, the scalar Poincaré inequality.

Example 2.5 (Gaussians). It is well known that the OU process satisfies the Poincaré inequality with constant $\alpha = 1$. Thus, it satisfies the trace Poincaré inequality (2.5) with $\alpha = 1$. For an alternative proof, see [ABY19, Thm. 1.2].

2.6. **Subexponential concentration and expectation bounds.** We are now prepared to present our main result. It demands several hypotheses, which will be enforced throughout the paper.

Assumption 2.6 (Conditions). We assume that

- (1) The Markov process $(Z_t: t \ge 0) \subset \Omega$ is reversible and homogeneous, with initial value Z_0 and stationary measure μ .
- (2) The process admits a trace Poincaré inequality (2.5) with constant α . Equivalently, the process admits a scalar Poincaré inequality with the same constant α .
- (3) The class of valid functions is suitably restricted so that manipulations of expectations, limits, and derivatives are justified.

Under Assumption 2.6, we will deduce subexponential concentration of the random matrix f(Z) around its mean $\mathbb{E}_{u} f$, where we measure the size of deviations with the ℓ_2 operator norm $\|\cdot\|$.

Theorem 2.7 (Subexponential Concentration). *Enforce Assumption 2.6.* Let $f: \Omega \to \mathbb{H}_d$ be a matrix-valued function, and define the variance proxy

$$v_f \coloneqq \left\| \left\| \Gamma(f)(z) \right\| \right\|_{L_{\infty}(\mu)}.$$

For all $\lambda > 0$,

$$\mathbb{P}_{\mu} \{ \| \boldsymbol{f} - \mathbb{E}_{\mu} \boldsymbol{f} \| \ge \sqrt{\alpha \nu_{\boldsymbol{f}}} \cdot \lambda \} \le 6d \cdot e^{-\lambda}.$$
 (2.6)

In particular,

$$\mathbb{E}_{\mu} \| \boldsymbol{f} - \mathbb{E}_{\mu} \boldsymbol{f} \| \le \log(6ed) \cdot \sqrt{\alpha v_{\boldsymbol{f}}}. \tag{2.7}$$

The proof of Theorem 2.7 appears in Section 6 after we present some more background on matrix-valued Markov processes. Note that we have made no effort to refine constants.

The main point of the tail bound (2.6) is that the random matrix f(Z) exhibits exponential concentration on the scale $\sqrt{\alpha v_f}$. The variance proxy v_f is analogous to a global bound on the

Lipschitz constant of f. Be aware that we cannot achieve tail decay faster than exponential under the sole assumption of a Poincaré inequality, so this approach may not capture the strongest possible concentration. The leading constant in (2.6) reflects the ambient dimension d of the matrix; this feature is typical of matrix concentration bounds.

The expectation bound (2.7) shows that the average value of $\|\mathbf{f} - \mathbb{E}_{\mu} \mathbf{f}\|$ is proportional to the square root $\sqrt{v_f}$ of the variance proxy and to the logarithm of the ambient dimension. For many examples, the optimal bound contains the *square-root* of the logarithm, but the result (2.7) is nontrivial and informative.

Example 2.8 (Gaussian Series). Theorem 2.7 applies to a matrix-valued function of a standard normal vector. For instance, according to (2.7), a matrix Gaussian series satisfies the expectation bound

$$\mathbb{E}\left\|\sum_{i=1}^{n} X_i A_i\right\| \leq \log(6ed) \cdot \left\|\sum_{i=1}^{n} A_i^2\right\|^{1/2}.$$

Modulo the constant and the power on the logarithm, this bound is qualitatively correct for worst-case examples. On the other hand, the subexponential tail bound (2.6) does not reproduce the actual subgaussian behavior. See [Tro15, Chap. 5] for discussion. Example 7.3 describes an application to Gaussian chaos that requires tools that are more delicate than Theorem 2.7.

Remark 2.9 (Extensions). The bounds in Theorem 2.7 can be refined in several ways. We can replace the variance proxy v_f with less stringent measures of the size of the carré du champ $\Gamma(f)$, such as the expected Schatten q-norm with $q \approx \log d$. It is also possible to replace the ambient dimension d with a measure of the intrinsic dimension of the random matrix f(Z). See Section 7.

Remark 2.10 (Rectangular case). By a standard formal argument, we can extend all the results here to a function $h: \Omega \to \mathbb{M}_{d_1 \times d_2}$ that takes values in the $d_1 \times d_2$ complex matrices. To do so, we simply apply our results to the self-adjoint function

$$f(z) = \begin{bmatrix} 0 & h(z) \\ h(z)^* & 0 \end{bmatrix} \in \mathbb{H}_{d_1+d_2} \quad \text{for } z \in \Omega.$$

See [Tro15, Sec. 2.1.17] for details.

2.7. **Examples.** To indicate the scope of Theorem 2.7, let us present some more examples. Most of these examples are actually known to exhibit subgaussian matrix concentration, but there is at least one case (Section 2.7.4) where the results here are currently the best available.

2.7.1. Log-concave measures. The Gaussian case is a particular example of a more general result for log-concave measures. Suppose that $J: \mathbb{R}^n \to \mathbb{R}$ is a strongly convex function that satisfies Hess $J \geqslant \eta \mathbf{I}$ uniformly. Construct the probability measure μ on \mathbb{R}^n whose density is proportional to e^{-J} . In this example, we briefly discuss concentration of matrix-valued functions f(X) where $X \sim \mu$. This model is interesting because it captures a type of negative dependence.

The appropriate Markov process $(X_t : t \ge 0)$ evolves with the stochastic differential equation

$$dX_t = -\nabla J(X_t) dt + \sqrt{2} dB_t$$
 with initial value $X_0 \in \mathbb{R}^n$,

where $(\mathbf{B}_t : t \ge 0) \subset \mathbb{R}^n$ is Brownian motion. The stationary distribution is μ , and the matrix carré du champ is

$$\Gamma(\mathbf{f})(\mathbf{x}) = \sum_{i=1}^{n} (\partial_i \mathbf{f}(\mathbf{x}))^2 \text{ for } \mathbf{x} \in \mathbb{R}^n.$$

It is well known that these diffusions satisfy a Poincaré inequality with constant $\alpha = 1/\eta$; see [BGL14, Cor. 4.8.2]. Therefore, Theorem 2.7 applies.

A fortiori, these log-concave measures also satisfy a local Poincaré inequality, which leads to subgaussian matrix concentration [HT20, Sec. 2.12.2].

2.7.2. Riemannian manifolds with positive curvature. More generally, let (M,\mathfrak{g}) be a compact Riemannian manifold with co-metric \mathfrak{g} . The manifold carries a canonical Riemannian probability measure $\mu_{\mathfrak{g}}$. The diffusion whose infinitesimal generator is the Laplace–Beltrami operator $\Delta_{\mathfrak{g}}$ is called the Brownian motion on the manifold. This is a reversible, ergodic Markov process. Its matrix carré du champ takes the form

$$\Gamma(\mathbf{f})(z) = \sum_{i,j} g^{ij}(z) \left(\partial_i \mathbf{f}(z)\right) \left(\partial_j \mathbf{f}(z)\right) \quad \text{for } \mathbf{f} : \Omega \to \mathbb{H}_d.$$
 (2.8)

The co-metric g and the partial derivatives $\partial_i f$ are computed with respect to local coordinates. See [BGL14] for an introduction to diffusions on manifolds; the companion paper [HT20] treats matrix-valued functions on manifolds.

Suppose that the manifold has uniformly positive Ricci curvature, where the Ricci tensor has eigenvalues bounded below by ρ . By now, it is a classic fact that the Brownian motion on this manifold satisfies the scalar Poincaré inequality with constant $\alpha = \rho^{-1}$. See [BGL14, Sec. 4.8].

Example 2.11 (Sphere). For $n \ge 2$, the unit sphere $\mathbb{S}^n \subset \mathbb{R}^{n+1}$ is a Riemannian submanifold of \mathbb{R}^{n+1} . Its canonical measure is the uniform probability distribution, and the carré du champ of the Brownian motion on the sphere is computed using (2.8). The sphere has positive Ricci curvature with $\rho = n - 1$, so it admits a Poincaré inequality with $\alpha = (n-1)^{-1}$. Thus, matrix-valued functions on the sphere satisfy exponential matrix concentration.

A similar story can be told about every positively curved manifold. In fact, in this setting, we even have subgaussian matrix concentration because of the stronger arguments in [HT20].

2.7.3. *Products*. Consider a probability space (Ω, μ) . It is common to work with multivariate functions defined on the product space $(\Omega^n, \mu^{\otimes n})$. There is a standard construction [VH16, Sec. 2.3.2] of a Markov process on the product space. Aoun et al. [ABY19] verify that the (matrix) carré du champ of this process is

$$\Gamma(f)(z) = \frac{1}{2} \sum_{i=1}^{n} \mathbb{E}_{Z \sim \mu} \left[\left(f(z_1, \dots, z_n) - f(z_1, \dots, z_{i-1}, Z, z_{i+1}, \dots, z_n) \right)^2 \right].$$

This is the sum of squared discrete derivatives, each averaging over perturbations in a single coordinate. The variance proxy v_f takes the form

$$v_{\mathbf{f}} = \operatorname{ess \, sup}_{\mathbf{z} \in \Omega^{n}} \frac{1}{2} \left\| \sum_{i=1}^{n} \mathbb{E}_{Z \sim \mu} \left[\left(\mathbf{f}(z_{1}, \dots, z_{n}) - \mathbf{f}(z_{1}, \dots, z_{i-1}, Z, z_{i+1}, \dots, z_{n}) \right)^{2} \right] \right\|$$

The variance proxy coincides with the matrix bounded difference that arises in Paulin et al. [PMT16]. Aoun et al. prove that the Markov process satisfies a trace Poincaré inequality (2.5) with constant $\alpha = 1$. Therefore, Theorem 2.7 yields a suboptimal version of the exponential matrix Efron–Stein inequality [PMT16, Thm. 4.3]. See [ABY19, HT20] for more details.

Remark 2.12 (Bernstein concentration?). In the scalar case, Bobkov & Ledoux [BL97, Cor. 3.2] have shown that a function $f:\Omega^n\to\mathbb{R}$ on a product space $(\Omega^n,\mu^{\otimes n})$ exhibits Bernstein-type concentration when μ admits a Poincaré inequality with constant α . In detail,

$$\mathbb{P}\left\{|f - \mathbb{E}\,f| > t\right\} \leq \exp\left(\frac{-\mathsf{c} t^2}{\nu + Bt}\right) \quad \text{with} \quad \nu = \left\|\sum_{i=1}^n (\partial_i f)^2\right\|_{L_\infty} \quad \text{and} \quad B = \max_i \|\partial_i f\|_{L_\infty}.$$

The constant c depends on the constant α in the Poincaré inequality. Our approach does not seem powerful enough to transfer this insight to the matrix setting. Nevertheless, our paper [HT20] demonstrates that a *local* Poincaré inequality is sufficient to achieve Bernstein concentration.

2.7.4. Stochastic covering property. Aoun et al. [ABY19] have considered a model for negatively dependent functions on the hypercube $\{0,1\}^n$, namely the class of measures with the stochastic covering property (SCP). For a k-homogeneous measure μ with the SCP, it is possible to construct a Markov

process that satisfies the trace Poincaré inequality (2.5) with constant 2k. Thus, Theorem 2.7 applies. See [PP14, HS19, ABY19] for a more complete discussion of this example.

Remark 2.13 (Subgaussian concentration?). Although the Markov process associated with an SCP measure satisfies a log-Sobolev inequality, we do not know if it satisfies the local Poincaré inequality that we would need to activate the subgaussian concentration inequalities in [HT20].

2.7.5. Expander graphs. Let $G=(\Omega,E)$ be a k-regular, connected, undirected graph. We can construct a Markov process $(Z_t:t\geq 0)\subset \Omega$ by taking a continuous-time random walk on the vertex set Ω . The stationary measure μ of the random walk is the uniform measure on Ω . The carré du champ operator takes the form

$$\Gamma(\mathbf{f})(z) = \frac{1}{2k} \sum_{(z',z) \in E} (\mathbf{f}(z') - \mathbf{f}(z))^{2}.$$

In other words, the "squared gradient" is just the half the average squared difference between the matrix at the current vertex and its k neighbors.

$$v_f := \max_{z \in \Omega} \left\| \frac{1}{2k} \sum_{(z',z) \in E} (f(z') - f(z))^2 \right\|.$$

The variance proxy is just the maximum "squared gradient" at any vertex.

Assume that the Markov process satisfies the (scalar) Poincaré inequality, Proposition 2.4(1), with constant α . In this case, we say that G is an α -expander graph. According to Theorem 2.7, a matrix-valued function $\mathbf{f}:\Omega\to\mathbb{H}_d$ on an α -expander graph satisfies a subexponential matrix concentration inequality:

$$\mathbb{P}\left\{\|\boldsymbol{f} - \mathbb{E}_{\mu}\boldsymbol{f}\| \geq \sqrt{\alpha v_{\boldsymbol{f}}} \cdot \lambda\right\} \leq 6d \cdot \mathrm{e}^{-\lambda}.$$

Therefore, local control over the fluctuations yields global concentration around the mean. The number of vertices where the function departs from its mean is controlled by the quality α of the expander.

This example is potentially interesting because the Markov process does not satisfy a dimension-independent (modified) log-Sobolev inequality [BTo6, Sec. 5]. Indeed, to achieve the functional inequality

$$\operatorname{Ent}_{\mu}(f) \leq \beta \cdot \mathscr{E}(f, \log f)$$
 for all $f: \Omega \to \mathbb{R}$,

it is necessary that $\beta \ge \text{const} \cdot \log(\#\Omega)$. See [Mün19, Sec. 4.3] for related results on curvature-dimension conditions of Bakry–Émery type.

Remark 2.14 (Matrix Expander Chernoff). Although the modified log-Sobolev inequality fails, it is still possible to establish subgaussian concentration for the ergodic averages of a matrix-valued random walk on an expander graph [GLSS18].

3. Related work

3.1. **Markov processes.** Much of the classical research on Markov processes concerns the relationship between the geometry of the state space and the behavior of canonical diffusion processes (e.g., Brownian motion on a Riemannian manifold). For an introduction, we recommend the lecture notes [VH16]. A more comprehensive source is the treatise [BGL14].

Matrix-valued Markov processes first arose in the mathematical physics literature as a model for the evolution of a quantum system. Some of the foundational works include Davies [Dav69] and Lindblad [Lin76]. Quantum information theory has provided a new impetus for studying matrix-valued Markov processes; see [KT13] for a discussion and some background references.

Here, we are interested in a mixed classical-quantum setting, where a classical Markov process drives a matrix-valued function. Surprisingly, this model does not seem to have received much

attention until the last few years. See Cheng et al. [CHT17] for a more expansive framework that includes this case. Other foundational results appear in [ABY19, HT20].

3.2. **Functional inequalities.** In the scalar setting, the connection between functional inequalities, convergence of Markov processes, and concentration is a long-standing topic of research. References include [Ledo1, BLM13, BGL14, VH16].

Functional inequalities for matrices were originally formulated in the mathematical physics literature; for example, see the work of Gross [Gro75]. The application of functional inequalities to the ergodicity of matrix-valued Markov processes dates back at least as far as the papers [MOZ98, OZ99].

Functional inequalities in the mixed classical-quantum setting seem to have a more recent vintage. Chen & Tropp [CT14] formulated subadditivity properties for tracial entropy-like quantities, including the trace variance (2.4). They showed that these properties imply some Sobolev and modified log-Sobolev-type inequalities for random matrices, and they obtained some restricted matrix concentration inequalities. Some of the partial results from [CT14] were completed in [HZ15, PV15, CH16].

Cheng et al. [CHT17] developed a framework for studying Markov processes in the mixed classical-quantum setting (and beyond), and they showed an equivalence between tracial log-Sobolev inequalities and exponential ergodicity of the trace entropy. Further results and implications for concentration appear in [CH19]. At present, we do not have a full picture of the relationships between matrix functional inequalities and matrix concentration.

Van Handel (personal communication) has pointed out that we can derive nonlinear matrix concentration for functions of a log-concave measure with a strongly convex potential by combining Pisier's method [Pis86], the (sharp) noncommutative Khintchine inequality [Buco1, Tro16], and Caffarelli's contraction theorem [Cafoo]. This approach gives subgaussian concentration, which is better than we can obtain via the trace Poincaré inequality for log-concave measures, but it apparently does not extend beyond this setting.

3.3. From Poincaré to concentration. It has been recognized for about 40 years that Poincaré inequalities imply exponential concentration. Gromov & Milman [GM83, Thm. 4.1] prove such a theorem in the context of Riemannian manifolds. The standard argument, a recursive estimate of the moment generating function, is attributed to Aida & Stroock [AS94]. For a textbook presentation, see [BLM13, Sec. 3.6].

Consider a matrix Poincaré inequality of the form

$$Var_{\mu}[f] \leq \alpha \cdot \mathscr{E}(f). \tag{3.1}$$

The argument in Proposition 2.4 shows that this matrix Poincaré inequality (3.1) is also equivalent to a scalar Poincaré inequality with the same constant α . The papers [CH16, CHT17, CH19] demonstrate that (3.1) leads to some inequalities for the matrix variance (2.4) and its trace, but these approaches do not lead to matrix concentration inequalities like Theorem 2.7.

Aoun, Banna, and Youssef [ABY19] have recently shown that the matrix Poincaré inequality (3.1) does imply exponential concentration of a random matrix about its mean with respect to the ℓ_2 operator norm. Modulo constants, their result is equivalent with the tail bound (2.6), but it is weaker than the bounds in Theorem 7.1. The proof in [ABY19] is a direct analog of the argument of Aida & Stroock [AS94]. But, in the matrix setting, the recursive estimate requires some heavy lifting. Another contribution of the paper [ABY19] is to establish that some particular matrix-valued Markov processes satisfy (3.1). Nevertheless, Proposition 2.4 indicates that no additional effort is required for this end.

Our approach is similar in spirit to the work of Aoun et al. [ABY19], but we use a symmetrization argument to avoid the difficult recursion. For related work in the scalar setting, see [BL97, Sec. 4]. In a companion paper [HT20], we show that *local* Poincaré inequalities lead to much stronger ergodicity and concentration properties. The theory in the companion paper is significantly more involved than

the development here, so we have chosen to separate them. For results on (local) Poincaré inequalities in noncommutative probability spaces, see Junge & Zeng [JZ15].

4. Subadditivity

To control the variance of a function of several independent variables, it is helpful to understand the influence of each individual variable. As in the scalar setting, the matrix variance can be bounded by a sum of conditional variances. We can control each conditional variance by a conditional application of a trace Poincaré inequality. For simplicity, we focus on the case that is relevant to our proof, but these results hold in greater generality (products of more than two spaces that may carry different measures). Some of the material in this section is drawn from [CT14, CH16]. See [Led97] for some of the classic work on subadditivity and concentration.

4.1. **Influence of a coordinate.** Consider the product space $(\Omega^2, \mu \otimes \mu)$. We want to study how individual coordinates affect the behavior of a matrix-valued function $\mathbf{g}: \Omega^2 \to \mathbb{H}_d$.

First, introduce notation for the expectation of the function with respect to each coordinate:

$$\mathbb{E}_1[\boldsymbol{g}](z_2) := \mathbb{E}_{Z \sim \mu}[\boldsymbol{g}(Z, z_2)] \in \mathbb{H}_d \quad \text{for all } z_1 \in \Omega;$$

$$\mathbb{E}_2[\boldsymbol{g}](z_1) := \mathbb{E}_{Z \sim \mu}[\boldsymbol{g}(z_1, Z)] \in \mathbb{H}_d \quad \text{for all } z_2 \in \Omega.$$

The coordinate-wise variance is the positive-semidefinite random matrix

$$\operatorname{Var}_{i}[\boldsymbol{g}] := \mathbb{E}_{i} \left[(\boldsymbol{g} - \mathbb{E}_{i} \, \boldsymbol{g})^{2} \right] \in \mathbb{H}_{d}^{+} \text{ for } i = 1, 2.$$

This matrix reflects the fluctuation in the *i*th coordinate, with the other coordinate held fixed. Similarly, we can introduce the coordinate-wise carré du champ operator and Dirichlet form:

$$\Gamma_{1}(\mathbf{g})(z_{1}, z_{2}) := \lim_{t \downarrow 0} \frac{1}{2t} \mathbb{E}\left[\left(\mathbf{g}(Z_{t}, z_{2}) - \mathbf{g}(Z_{0}, z_{2})\right)^{2} \middle| Z_{0} = z_{1}\right] \in \mathbb{H}_{d}^{+};$$

$$\mathscr{E}_{1}(\mathbf{g})(z_{2}) := \lim_{t \downarrow 0} \frac{1}{2t} \mathbb{E}_{Z \sim \mu}\left[\left(\mathbf{g}(Z_{t}, z_{2}) - \mathbf{g}(Z_{0}, z_{2})\right)^{2} \middle| Z_{0} = Z\right] \in \mathbb{H}_{d}^{+}.$$
(4.1)

As usual, $(Z_t : t \ge 0)$ is a realization of the Markov process with initial value Z_0 . We make analogous definitions for the second coordinate i = 2.

Last, we extend the carré du champ operator and the Dirichlet form to bivariate functions:

$$\Gamma(\mathbf{g}) := \Gamma_1(\mathbf{g}) + \Gamma_2(\mathbf{g}) \in \mathbb{H}_d^+; \tag{4.2}$$

$$\mathscr{E}(\mathbf{g}) := \mathbb{E}_{\mu \otimes \mu} \left[\mathscr{E}_1(\mathbf{g}) + \mathscr{E}_2(\mathbf{g}) \right] \in \mathbb{H}_d^+. \tag{4.3}$$

These formulas have a heuristic interpretation: the squared derivative of a bivariate function is the sum of the squared partial derivatives.

4.2. **Trace variance is subadditive.** Observe that the trace variance is controlled by the sum of the coordinate-wise variances.

Fact 4.1 (Trace variance: Subadditivity). Let $\mathbf{g}: \Omega^2 \to \mathbb{H}_d$ be a matrix-valued function on the product space $(\Omega^2, \mu \otimes \mu)$. Then

$$\operatorname{tr} \operatorname{Var}_{\mu \otimes \mu}[\boldsymbol{g}] \leq \mathbb{E}_2 \operatorname{tr} \operatorname{Var}_1[\boldsymbol{g}] + \mathbb{E}_1 \operatorname{tr} \operatorname{Var}_2[\boldsymbol{g}].$$

This result is due to Chen & Tropp [CT14], who showed that other matrix functions are also subadditive. Later, Cheng & Hsieh [CH16] noticed that an analogous result holds without the trace. Similar decompositions are also valid for functions on the n-fold product $(\Omega^n, \mu^{\otimes n})$.

Proof. The proof is the same as in the scalar setting. Writing $\mathbb{E} = \mathbb{E}_{\mu \otimes \mu}$ for the total expectation,

$$\mathbb{E} \operatorname{Var}_{\mu \otimes \mu}[\mathbf{g}] := \mathbb{E} \left[(\mathbf{g} - \mathbb{E} \, \mathbf{g})^2 \right] = \mathbb{E} \left[(\mathbf{g} - \mathbb{E}_1 \, \mathbf{g})^2 + (\mathbb{E}_1 \, \mathbf{g} - \mathbb{E}_1 \, \mathbb{E}_2 \, \mathbf{g})^2 \right]$$

$$\leq \mathbb{E}[(\boldsymbol{g} - \mathbb{E}_1 \, \boldsymbol{g})^2] + \mathbb{E}[(\boldsymbol{g} - \mathbb{E}_2 \, \boldsymbol{g})^2]$$

= $\mathbb{E}_2 \operatorname{Var}_1[\boldsymbol{g}] + \mathbb{E}_1 \operatorname{Var}_2[\boldsymbol{g}].$

The first line relies on orthogonality of the conditionally centered random matrices. The second line requires the operator convexity of the square, applied conditionally. Last, take the trace.

4.3. **Trace Poincaré inequalities are subadditive.** If the Markov process satisfies a trace Poincaré inequality, then the variance of a bivariate function also satisfies a trace Poincaré inequality.

Proposition 4.2 (Trace Poincaré: Subadditivity). Suppose that the Markov process satisfies a trace Poincaré inequality (2.5) with constant α . Let $\mathbf{g}:\Omega^2\to\mathbb{H}_d$ be a suitable bivariate matrix-valued function. Then

$$\operatorname{tr} \operatorname{Var}_{\mu \otimes \mu}[\mathbf{g}] \leq \alpha \cdot \operatorname{tr} \mathscr{C}(\mathbf{g}).$$

Proof. Start with Fact 4.1. Apply the trace Poincaré inequality (2.5) coordinate-wise to control each of the two coordinate-wise variances. Last, introduce the definition (4.3) of the Dirichlet form for a bivariate function.

5. Chain rule inequality for the Dirichlet form

The key new tool in our approach is a simple trace inequality for the matrix Dirichlet form that shows how it interacts with composition.

Proposition 5.1 (Chain rule inequality). *Enforce Assumption 2.6.* Let $\varphi : \mathbb{R} \to \mathbb{R}$ be a scalar function whose squared derivative $\psi = (\varphi')^2$ is convex. Then

$$\operatorname{tr} \mathscr{C}(\varphi(\boldsymbol{f})) = \mathbb{E}_{\mu} \operatorname{tr} \Gamma(\varphi(\boldsymbol{f})) \leq \mathbb{E}_{\mu} \operatorname{tr} \left[\Gamma(\boldsymbol{f}) \, \psi(\boldsymbol{f}) \right] \quad \text{for all suitable } \boldsymbol{f} : \Omega \to \mathbb{H}_d.$$

In particular,

$$\operatorname{tr} \mathscr{C}(\varphi(\boldsymbol{g})) = \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \Gamma(\varphi(\boldsymbol{g})) \leq \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \left[\Gamma(\boldsymbol{g}) \, \psi(\boldsymbol{g}) \right] \quad \textit{for all suitable } \boldsymbol{g} : \Omega^2 \to \mathbb{H}_d.$$

The proof of Proposition 5.1 consumes the rest of this section.

For context, recall that the carré du champ operator Γ of a (scalar-valued, reversible) diffusion process satisfies a chain rule [BGL14, Sec. 1.11]:

$$\Gamma(\varphi(f)) = \Gamma(f) \varphi'(f)^2$$
 for smooth f and φ .

Proposition 5.1 provides a substitute for this relation for an arbitrary reversible Markov process that takes matrix values. In exchange for the wider applicability, we need some additional averaging (provided by the Dirichlet form); we must restrict our attention to functions φ with a convexity property; and the equality is relaxed to an inequality. In the scalar case, Proposition 5.1 is related to the Stroock–Varopoulos inequality [Str84, Var85].

5.1. **Mean-value inequality for trace functions.** The argument hinges on an elementary trace inequality for deterministic matrices. This result is obtained by lifting a numerical inequality to self-adjoint matrices. A very similar statement [MJC⁺14, Lem. 3.4] animates the exchangeable pairs approach to matrix concentration, which is motivated by work in the scalar setting [Chao₇].

Lemma 5.2 (Mean-value trace inequality). Let $A, B \in \mathbb{H}_d$. Let $\varphi : \mathbb{R} \to \mathbb{R}$ be a scalar function whose squared derivative $\psi = (\varphi')^2$ is convex. Then

$$\operatorname{tr}\left[\left(\varphi(\boldsymbol{A})-\varphi(\boldsymbol{B})\right)^{2}\right] \leq \frac{1}{2}\operatorname{tr}\left[\left(\boldsymbol{A}-\boldsymbol{B}\right)^{2}\left(\psi(\boldsymbol{A})+\psi(\boldsymbol{B})\right)\right].$$

Proof. Let $a, b \in \mathbb{R}$. The fundamental theorem of calculus and Jensen's inequality together deliver the relations

$$(\varphi(a) - \varphi(b))^{2} = (a - b)^{2} \left[\int_{0}^{1} d\tau \ \varphi'(\tau a + (1 - \tau)b) \right]^{2}$$

$$\leq (a - b)^{2} \int_{0}^{1} d\tau \ \psi(\tau a + (1 - \tau)b)$$

$$\leq (a - b)^{2} \int_{0}^{1} d\tau \left[\tau \psi(a) + (1 - \tau)\psi(b) \right] = \frac{1}{2} (a - b)^{2} (\psi(a) + \psi(b)).$$

The generalized Klein inequality [Pet94, Prop. 3] allows us to lift this numerical fact to a trace inequality for matrices A, $B \in \mathbb{H}_d$.

5.2. **Proof of Proposition 5.1.** The result follows from a short calculation. First, we use the definition (2.3) of the Dirichlet form as a limit:

$$\operatorname{tr} \mathscr{E}(\varphi(\boldsymbol{f})) = \lim_{t \downarrow 0} \frac{1}{2t} \mathbb{E}_{Z \sim \mu} \left[\operatorname{tr} \left[\left(\varphi(\boldsymbol{f}(Z_t)) - \varphi(\boldsymbol{f}(Z_0)) \right)^2 \right] \middle| Z_0 = Z \right]$$

$$\leq \lim_{t \downarrow 0} \frac{1}{4t} \mathbb{E}_{Z \sim \mu} \operatorname{tr} \left[\left(\boldsymbol{f}(Z_t) - \boldsymbol{f}(Z_0) \right)^2 \left(\psi(\boldsymbol{f}(Z_t)) + \psi(\boldsymbol{f}(Z_0)) \right) \middle| Z_0 = Z \right]$$

$$= \lim_{t \downarrow 0} \frac{1}{2t} \mathbb{E}_{Z \sim \mu} \operatorname{tr} \left[\left(\boldsymbol{f}(Z_t) - \boldsymbol{f}(Z_0) \right)^2 \psi(\boldsymbol{f}(Z_0)) \middle| Z_0 = Z \right]$$

$$= \mathbb{E}_{Z \sim \mu} \operatorname{tr} \left[\lim_{t \downarrow 0} \frac{1}{2t} \mathbb{E} \left[\left(\boldsymbol{f}(Z_t) - \boldsymbol{f}(Z_0) \right)^2 \middle| Z_0 = Z \right] \psi(\boldsymbol{f}(Z)) \right]$$

$$= \mathbb{E}_{\mu} \operatorname{tr} \left[\Gamma(\boldsymbol{f}) \psi(\boldsymbol{f}) \right].$$

The inequality is Lemma 5.2. To reach the third line, we use the fact that (Z_0, Z_t) is an exchangeable pair for each $t \geq 0$, a consequence of the reversibility of the Markov process $(Z_t : t \geq 0)$ and the fact that $Z_0 \sim \mu$. Last, we condition on the value of Z_0 , invoke dominated convergence to pass the expectation through the limit, and we apply the definition (2.2) of the carré du champ operator.

6. Exponential moments

Our main technical result is a bound for the exponential moments of a general matrix-valued function on the state space. In contrast to the usual approach of bounding the moment generating function, we will compute the expectation of a hyperbolic function.

Theorem 6.1 (Exponential moments). *Enforce Assumption 2.6.* Let $\mathbf{f}: \Omega \to \mathbb{H}_d$ be a function with $\mathbb{E}_{\mu} \mathbf{f} = \mathbf{0}$. For $\theta > 0$,

$$\mathbb{E}_{\mu} \operatorname{tr} \cosh(\theta \mathbf{f}) \leq d \cdot \left[1 + \frac{\alpha \theta^{2} \operatorname{tr} \mathscr{E}(\mathbf{f})}{(1 - \alpha v_{\mathbf{f}} \theta^{2}/2)_{+}} \right]. \tag{6.1}$$

The variance proxy v_f is defined in (2.7), and $\bar{\text{tr}}$ is the normalized trace.

The proof of Theorem 6.1 occupies the rest of the section. But first, we use this moment bound to derive our main result, Theorem 2.7.

Proof of Theorem 2.7 from Theorem 6.1. Without loss, assume that $\mathbb{E}_{\mu} f = 0$. We use the matrix moment method [AWo2, Oli10]:

$$\mathbb{P}_{\mu}\left\{\left\|\boldsymbol{f}\right\| \geq \lambda\right\} \leq \inf_{\theta > 0} \frac{1}{\cosh(\theta\lambda)} \cdot \mathbb{E}_{\mu} \cosh(\theta \left\|\boldsymbol{f}\right\|) = \inf_{\theta > 0} \frac{2}{\mathrm{e}^{\theta\lambda} + \mathrm{e}^{-\theta\lambda}} \cdot \mathbb{E}_{\mu} \left\|\cosh(\theta\boldsymbol{f})\right\|$$

$$\leq \inf_{\theta>0} \frac{2}{\mathrm{e}^{\theta\lambda}} \cdot \mathbb{E}_{\mu} \operatorname{tr} \cosh(\theta \boldsymbol{f}) \leq 2d \cdot \inf_{\theta>0} \mathrm{e}^{-\theta\lambda} \cdot \left[1 + \frac{\alpha \theta^{2} \operatorname{tr} \boldsymbol{\mathcal{E}}(\boldsymbol{f})}{(1 - \alpha v_{\boldsymbol{f}} \theta^{2}/2)_{+}} \right]$$

$$\leq 2d \cdot \mathrm{e}^{-\lambda/\sqrt{\alpha v_{\boldsymbol{f}}}} \cdot \left[1 + \frac{2 \operatorname{tr} \boldsymbol{\mathcal{E}}(\boldsymbol{f})}{v_{\boldsymbol{f}}} \right] \leq 6d \cdot \mathrm{e}^{-\lambda/\sqrt{\alpha v_{\boldsymbol{f}}}}.$$

The first inequality is Markov's. The second relation is the spectral mapping theorem. The ℓ_2 operator norm of a positive-definite matrix is obviously bounded by its trace. Then invoke Theorem 6.1 to control the moment. We have chosen $\theta = (\alpha v_f)^{-1/2}$. Last, we have noted that $\operatorname{tr} \mathscr{C}(f) \leq \|\mathscr{C}(f)\| \leq v_f$. To finish the proof of the tail bound (2.6), make the change of variables $\lambda \mapsto \lambda \sqrt{\alpha v_f}$.

The expectation bound (2.7) follows when we integrate the tail bound (2.6), taking into account that the probability is also bounded by one; we omit the details.

- 6.1. **Proof of Theorem 6.1.** Our goal is to develop an exponential moment bound for a function $f: \Omega \to \mathbb{H}_d$ that satisfies $\mathbb{E}_{\mu} f = 0$. We will need to work with both the hyperbolic sine and cosine, passing between them using simple identities. After writing this paper, we learned that this proof is a matrix analog of an argument proposed by Bobkov & Ledoux [BL97, Sec. 4].
- 6.1.1. Symmetrization. The first step is to symmetrize the function. Let $Z, Z' \in \Omega$ be independent random variables, each with distribution μ . Since f(Z') has mean zero, a conditional application of Jensen's inequality yields

$$\mathbb{E}\operatorname{tr}\sinh^{2}(\theta \boldsymbol{f}(Z)) \leq \mathbb{E}\operatorname{tr}\sinh^{2}(\theta(\boldsymbol{f}(Z) - \boldsymbol{f}(Z'))) =: \mathbb{E}\operatorname{tr}\sinh^{2}(\theta \boldsymbol{g}(Z, Z')). \tag{6.2}$$

Indeed, since \sinh^2 is convex, the function $\operatorname{tr} \sinh^2$ is also convex [Pet94, Prop. 2]. We have defined the antisymmetric function $\mathbf{g}(z,z')=\mathbf{f}(z)-\mathbf{f}(z')$ for $z,z'\in\Omega$.

6.1.2. From moments to variance. Next, we will write the expectation as a variance. Consider the odd function $\varphi(s) = \sinh(\theta s)$. First, we claim that

$$\mathbb{E}\sinh(\theta \mathbf{g}(Z,Z'))=\mathbf{0}.$$

Indeed, using the antisymmetry of g and the oddness of φ ,

$$\mathbb{E} \sinh(\theta \mathbf{g}(Z, Z')) = \mathbb{E} \sinh(-\theta \mathbf{g}(Z', Z)) = -\mathbb{E} \sinh(\theta \mathbf{g}(Z', Z)) = -\mathbb{E} \sinh(\theta \mathbf{g}(Z, Z')).$$

The last identity holds because (Z, Z') is exchangeable.

As an immediate consequence,

$$\mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \sinh^{2}(\theta \boldsymbol{g}) = \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \sinh^{2}(\theta \boldsymbol{g}) - \operatorname{tr} \left[\left(\mathbb{E}_{\mu \otimes \mu} \sinh(\theta \boldsymbol{g}) \right)^{2} \right] = \operatorname{tr} \operatorname{Var}_{\mu \otimes \mu} \left[\sinh(\theta \boldsymbol{g}) \right].$$

The appearance of the variance gives us access to Poincaré inequalities.

6.1.3. *Poincaré inequality*. To continue, we apply Proposition 4.2, the trace Poincaré inequality for bivariate functions:

$$\mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \sinh^{2}(\theta \mathbf{g}) \leq \alpha \cdot \operatorname{tr} \mathscr{E}(\sinh(\theta \mathbf{g})) \leq \alpha \theta^{2} \cdot \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \left[\mathbf{\Gamma}(\mathbf{g}) \cosh^{2}(\theta \mathbf{g}) \right]. \tag{6.3}$$

The second inequality is the chain rule, Proposition 5.1, for the Dirichlet form. To activate it, we note that the squared derivative of $\varphi(s) = \sinh(\theta s)$ is the convex function $\psi(s) = \theta^2 \cosh^2(\theta s)$.

6.1.4. *Moment comparison*. A moment comparison argument allows us to isolate the exponential moment. Define the variance proxy of the bivariate function:

$$\nu_{\mathbf{g}} := \|\|\Gamma(\mathbf{g})\|\|_{L_{\infty}(\mu \otimes \mu)}. \tag{6.4}$$

Continuing from (6.3), the identity $\cosh^2 = 1 + \sinh^2$ implies that

$$\mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \sinh^2(\theta \boldsymbol{g}) \leq \alpha \theta^2 \cdot \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \boldsymbol{\Gamma}(\boldsymbol{g}) + \alpha \theta^2 \cdot \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \left[\boldsymbol{\Gamma}(\boldsymbol{g}) \sinh^2(\theta \boldsymbol{g}) \right]$$

$$\leq \alpha \theta^2 \cdot \operatorname{tr} \mathscr{C}(\boldsymbol{g}) + \alpha \theta^2 v_{\boldsymbol{g}} \cdot \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \sinh^2(\theta \boldsymbol{g}).$$

The second inequality is just the usual operator-norm bound for the trace. Rearrange this identity to arrive at

$$\mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \sinh^{2}(\theta \mathbf{g}) \leq \frac{\alpha \theta^{2} \operatorname{tr} \mathscr{E}(\mathbf{g})}{(1 - \alpha \nu_{\mathbf{g}} \theta^{2})_{+}}.$$
(6.5)

It remains to revert to the original function f.

6.1.5. Comparison of carré du champs. Let us compute the Dirichlet form $\mathcal{E}(g)$ and the variance proxy v_f in terms of the original function f. To that end, observe that the coordinate-wise carré du champ (4.1) satisfies

$$\Gamma_{1}(\boldsymbol{g})(z,z') = \mathbb{E}\lim_{t\downarrow 0} \frac{1}{2t} \left[\left(\boldsymbol{g}(Z_{t},z') - \boldsymbol{g}(Z_{0},z') \right)^{2} \middle| Z_{0} = z \right]$$

$$= \mathbb{E}\lim_{t\downarrow 0} \frac{1}{2t} \left[\left(\boldsymbol{f}(Z_{t}) - \boldsymbol{f}(Z_{0}) \right)^{2} \middle| Z_{0} = z \right] = \Gamma(\boldsymbol{f})(z).$$

A similar calculation reveals that $\Gamma_2(\mathbf{g})(z,z') = \Gamma(\mathbf{f})(z')$. Thus, the bivariate carré du champ (4.2) takes the form

$$\Gamma(\mathbf{g})(z,z') = \Gamma_1(\mathbf{g})(z,z') + \Gamma_2(\mathbf{g})(z,z') = \Gamma(\mathbf{f})(z) + \Gamma(\mathbf{f})(z'). \tag{6.6}$$

As a consequence, the Dirichlet form can be calculated as

$$\mathscr{E}(\mathbf{g}) = \mathbb{E}_{\mu \otimes \mu} \, \Gamma(\mathbf{g}) = \mathbb{E}_{\mu \otimes \mu} \left[\Gamma(\mathbf{f})(z) + \Gamma(\mathbf{f})(z') \right] = 2\mathscr{E}(\mathbf{f}). \tag{6.7}$$

The variance proxy (6.4) of the bivariate function satisfies

$$v_{\mathbf{g}} = \|\|\mathbf{\Gamma}(\mathbf{g})(z, z')\|\|_{L_{\infty}(\mu \otimes \mu)} \le \|\|\mathbf{\Gamma}(\mathbf{f})(z)\|\|_{L_{\infty}(\mu)} + \|\|\mathbf{\Gamma}(\mathbf{f})(z')\|\|_{L_{\infty}(\mu)} = 2v_{\mathbf{f}}.$$
(6.8)

The last relation is the definition (2.7) of the variance proxy v_f .

6.1.6. Endgame. Combining (6.2) and (6.5), we see that

$$\mathbb{E}_{\mu} \operatorname{tr} \sinh^{2}(\theta \boldsymbol{f}) \leq \frac{\alpha \theta^{2} \operatorname{tr} \boldsymbol{\mathcal{E}}(\boldsymbol{g})}{(1 - \alpha \nu_{\boldsymbol{g}} \theta^{2})_{+}} \leq \frac{2\alpha \theta^{2} \operatorname{tr} \boldsymbol{\mathcal{E}}(\boldsymbol{f})}{(1 - 2\alpha \nu_{\boldsymbol{f}} \theta^{2})_{+}}.$$

We have also used the relations (6.7) and (6.8) from the last section.

To compete the proof of (6.1), invoke the identity $\sinh^2(s) = (\cosh(2s) - 1)/2$ to see that

$$\mathbb{E}_{\mu}\operatorname{tr}\cosh(2\theta \boldsymbol{f}) \leq d + \frac{4\alpha\theta^{2}\operatorname{tr}\boldsymbol{\mathcal{E}}(\boldsymbol{f})}{(1 - 2\alpha\nu_{\boldsymbol{f}}\theta^{2})_{+}}.$$

Finally, introduce the normalized trace, \bar{tr} , and make the change of variables $\theta \mapsto \theta/2$ to arrive at

$$\mathbb{E}_{\mu}\operatorname{tr}\cosh(\theta \boldsymbol{f}) \leq d \cdot \left[1 + \frac{\alpha \theta^2 \operatorname{tr} \boldsymbol{\mathcal{E}}(\boldsymbol{f})}{(1 - \alpha \nu_{\boldsymbol{f}} \theta^2 / 2)_+}\right].$$

This is the stated result.

7. POLYNOMIAL MOMENTS

By a simple variation on the proof of Theorem 6.1, we can also obtain bounds for the polynomial moments of a random matrix.

Theorem 7.1 (Polynomial moments). *Enforce Assumption 2.6.* Let $\mathbf{f}: \Omega \to \mathbb{H}_d$ be a function with $\mathbb{E}_{\mu} \mathbf{f} = \mathbf{0}$. For q = 1 and $q \geq 1.5$,

$$\left(\mathbb{E}_{\mu}\operatorname{tr}\left|\boldsymbol{f}\right|^{2q}\right)^{1/(2q)} \leq \sqrt{2\alpha q^{2}} \cdot \left(\mathbb{E}_{\mu}\operatorname{tr}\boldsymbol{\Gamma}(\boldsymbol{f})^{q}\right)^{1/(2q)}.$$
(7.1)

By combining Theorem 7.1 with the moment method, we can obtain probability bounds for ||f||. Let us summarize how these results compare with the main result, Theorem 2.7. Observe that Theorem 7.1 gives a bound on the Schatten 2q-norm of the random matrix f(Z) in terms of the Schatten 2q-norm of $\Gamma(f)^{1/2}$. We have the relation

$$\left(\mathbb{E}_{\mu} \operatorname{tr} \mathbf{\Gamma}(\mathbf{f})^{q} \right)^{1/(2q)} \le d^{1/(2q)} \cdot \left\| \left\| \mathbf{\Gamma}(\mathbf{f})(z) \right\| \right\|_{L_{\infty}(\mu)}^{1/2} = d^{1/(2q)} \cdot \sqrt{v_{\mathbf{f}}}. \tag{7.2}$$

Therefore, Theorem 7.1 potentially yields stronger bounds than Theorem 6.1.

In particular, Theorem 7.1 applies even when $\Gamma(f)$ is not uniformly bounded. Example 7.3 illustrates why this flexibility is valuable. In Section 7.2, we show that slightly better polynomial moment bounds are possible when $\Gamma(f)$ is uniformly bounded.

Remark 7.2 (Missing Parameters). Theorem 7.1 also holds for $q \in (1, 1.5)$, with an extra factor of $\sqrt{2}$ on the right-hand side. The proof uses a variant of Proposition 5.1 that only requires the function ψ to be monotone.

Example 7.3 (Gaussian Chaos). Consider the matrix Gaussian chaos

$$f(X) = \sum_{i,j=1}^{n} X_i X_j A_{ij}$$
 where $X \sim \gamma_n$ and $A_{ij} = A_{ji} \in \mathbb{H}_d$.

To bound the trace moments of f(X), observe that the carré du champ takes the form

$$\Gamma(\mathbf{f})(\mathbf{x}) = \sum_{i=1}^{n} (\partial_i \mathbf{f}(\mathbf{x}))^2 = 4 \sum_{i=1}^{n} \left(\sum_{j=1}^{n} x_j \mathbf{A}_{ij} \right)^2.$$

Evidently, $\Gamma(f)$ is unbounded, so Theorem 2.7 does not apply. But Theorem 7.1 yields

$$\left(\mathbb{E}_{\gamma_n}\operatorname{tr}\left|\boldsymbol{f}\right|^{2q}\right)^{1/(2q)} \leq \sqrt{8q^2} \cdot \left(\mathbb{E}_{\gamma_n}\operatorname{tr}\left[\sum_{i=1}^n \left(\sum_{j=1}^n X_j \boldsymbol{A}_{ij}\right)^2\right]^q\right)^{1/(2q)}.$$
 (7.3)

We have used the fact that the Poincaré constant of the OU process is $\alpha = 1$. Further bounds can be obtained by applying Theorem 7.1 repeatedly. In the matrix setting, there are obstacles that prevent us from simplifying (7.3) completely (related to the fact that the partial transpose operator is not completely bounded).

In the scalar case d = 1, we can obtain more transparent conclusions. Consider the scalar-valued Gaussian chaos

$$f(\mathbf{X}) = \sum_{i,j=1}^{n} X_i X_j a_{ij}.$$

The most satisfying outcome takes place when $A = [a_{ij}]$ is positive semidefinite. In this case, the result (7.3) implies that

$$\left(\mathbb{E}_{\gamma_{n}} |f|^{2q}\right)^{1/(2q)} \leq \sqrt{8q^{2}} \cdot \left(\mathbb{E}_{\gamma_{n}} \left[\sum_{i=1}^{n} \left(\sum_{j=1}^{n} X_{j} a_{ij}\right)^{2}\right]^{q}\right)^{1/(2q)} \\
= \sqrt{8q^{2}} \cdot \left(\mathbb{E}_{\gamma_{n}} \left[\sum_{j,k=1}^{n} X_{j} X_{k} (A^{2})_{jk}\right]^{q}\right)^{1/(2q)} \\
\leq \sqrt{8q^{2} ||A||} \cdot \left(\mathbb{E}_{\gamma_{n}} |f|^{q}\right)^{1/2q} \leq \sqrt{8q^{2} ||A||} \cdot \left(\mathbb{E}_{\gamma_{n}} |f|^{2q}\right)^{1/4q}.$$

Solving, we obtain the bound

$$(\mathbb{E}_{\gamma_n} |f|^{2q})^{1/(2q)} \leq 8q^2 ||A||.$$

This result gives a suboptimal estimate for large moments of the Gaussian chaos; as $q \to \infty$, the moments should grow in proportion to $q \|A\|$ rather than $q^2 \|A\|$. For example, see [LT11, Cor. 3.9].

7.1. **Proof of Theorem 7.1.** For a parameter q=1 or $q\geq 1.5$, we wish to estimate the Schatten 2q-norm of a function $f:\Omega\to\mathbb{H}_d$ that satisfies $\mathbb{E}_{\mu}f=0$. The argument has the same structure as Theorem 6.1.

First, we symmetrize. Let $Z, Z' \in \Omega$ be independent random variables, each with distribution μ . Jensen's inequality implies that

$$\mathbb{E}\operatorname{tr}\left|\boldsymbol{f}\right|^{2q} \leq \mathbb{E}\operatorname{tr}\left|\boldsymbol{f}(Z) - \boldsymbol{f}(Z')\right|^{2q} =: \mathbb{E}\operatorname{tr}\left|\boldsymbol{g}(Z, Z')\right|^{2q}. \tag{7.4}$$

Since $|\cdot|^{2q}$ is convex, the function tr $|\cdot|^{2q}$ is also convex [Pet94, Prop. 2].

Define the signed moment function $\varphi(s) := \operatorname{sgn}(s) \cdot |s|^q$, which is odd. Note that its squared derivative $\psi(s) := (\varphi'(s))^2 = q^2 |s|^{2(q-1)}$ is convex. Since $\mathbf{g}(z,z') = \mathbf{f}(z) - \mathbf{f}(z')$ is antisymmetric,

$$\mathbb{E}_{\mu \otimes \mu} \operatorname{tr} |\boldsymbol{g}|^{2q} = \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \varphi(\boldsymbol{g})^{2}$$
$$= \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \varphi(\boldsymbol{g})^{2} - \operatorname{tr} \left[\left(\mathbb{E}_{\mu \otimes \mu} \varphi(\boldsymbol{g}) \right)^{2} \right] = \operatorname{tr} \operatorname{Var}_{\mu \otimes \mu} [\varphi(\boldsymbol{g})].$$

Apply the bivariate trace Poincaré inequality, Proposition 4.2:

$$\operatorname{tr} \operatorname{Var}_{\mu \otimes \mu} [\varphi(\mathbf{g})] \leq \alpha \cdot \operatorname{tr} \mathscr{E}(\varphi(\mathbf{g})) \leq \alpha q^2 \cdot \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \big[\Gamma(\mathbf{g}) \, \psi(\mathbf{g}) \big].$$

The second bound is the chain rule inequality, Proposition 5.1. In summary,

$$\mathbb{E}_{\mu \otimes \mu} \operatorname{tr} |\mathbf{g}|^{2q} \le \alpha q^2 \cdot \mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \left[\mathbf{\Gamma}(\mathbf{g}) |\mathbf{g}|^{2(q-1)} \right]. \tag{7.5}$$

This formula allow us to perform a moment comparison.

To isolate the carré du champ $\Gamma(g)$, invoke Hölder's inequality for the Schatten norms:

$$\mathbb{E}_{\mu \otimes \mu} \operatorname{tr} |\boldsymbol{g}|^{2q} \leq \alpha q^2 \left(\mathbb{E}_{\mu \otimes \mu} \operatorname{tr} \boldsymbol{\Gamma}(\boldsymbol{g})^q \right)^{1/q} \left(\mathbb{E}_{\mu \otimes \mu} \operatorname{tr} |\boldsymbol{g}|^{2q} \right)^{(q-1)/q}.$$

Rearrange the last display, and use the initial bound (7.4) to arrive at

$$\left(\left.\mathbb{E}_{\mu\otimes\mu}\operatorname{tr}\left|\boldsymbol{f}\right|^{2q}\right)^{1/(2q)}\leq\left(\left.\mathbb{E}_{\mu\otimes\mu}\operatorname{tr}\left|\boldsymbol{g}\right|^{2q}\right)^{1/(2q)}\leq\sqrt{\alpha q^{2}}\cdot\left(\left.\mathbb{E}_{\mu\otimes\mu}\operatorname{tr}\boldsymbol{\Gamma}(\boldsymbol{g})^{q}\right)^{1/(2q)}.$$

To finish the proof of (7.1), recall the expression (6.6) for the carré du champ $\Gamma(g)$. Thus,

$$\left(\mathbb{E}_{\mu\otimes\mu}\operatorname{tr}\mathbf{\Gamma}(\boldsymbol{g})^{q}\right)^{1/q} = \left(\mathbb{E}_{\mu\otimes\mu}\operatorname{tr}\left|\mathbf{\Gamma}(\boldsymbol{f})(z) + \mathbf{\Gamma}(\boldsymbol{f})(z')\right|^{q}\right)^{1/q} \leq 2\left(\mathbb{E}_{\mu}\operatorname{tr}\mathbf{\Gamma}(\boldsymbol{f})^{q}\right)^{1/q}.$$

This point follows from the triangle inequality. The argument is complete.

7.2. A variant of the argument. The intrinsic dimension of a positive-semidefinite matrix is

$$\operatorname{intdim}(A) := \frac{\operatorname{tr}(A)}{\|A\|} \quad \text{for } A \in \mathbb{H}_d^+.$$

We also set $\operatorname{intdim}(\mathbf{0}) = 0$. For a nonzero matrix A, the intrinsic dimension satisfies $1 \leq \operatorname{intdim}(A) \leq \operatorname{rank}(A)$. It can be interpreted as a continuous measure of the rank.

Suppose that q is a natural number. If we use the uniform bound (6.4) for the carré du champ instead of Hölder's inequality, we can apply the bound (7.5) iteratively to obtain

$$\mathbb{E}_{\mu \otimes \mu} \operatorname{tr} |\mathbf{g}|^{2q} \leq \operatorname{intdim}(\mathscr{E}(\mathbf{g})) \cdot \alpha^q q! \cdot v_{\mathbf{g}}^q.$$

The latter estimate improves over the uniform inequality that follows from Theorem 7.1 and (7.2).

ACKNOWLEDGMENTS

Ramon Van Handel offered valuable feedback on a preliminary version of this work, and we are grateful to him for the proof of Proposition 2.4. DH was funded by NSF grants DMS-1907977 and DMS-1912654. JAT gratefully acknowledges funding from ONR awards No0014-17-12146 and No0014-18-12363, and he would like to thank his family for their support in these difficult times.

REFERENCES

- [ABY19] R. Aoun, M. Banna, and P. Youssef. Matrix Poincaré inequalities and concentration, 2019.
- [AS94] S. Aida and D. Stroock. Moment estimates derived from Poincaré and logarithmic Sobolev inequalities. *Math. Res. Lett.*, 1(1):75–86, 1994.
- [AWo2] R. Ahlswede and A. Winter. Strong converse for identification via quantum channels. *IEEE Trans. Inform. Theory*, 48(3):569–579, 2002.
- [BGL14] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Cham, 2014.
- [BL97] S. Bobkov and M. Ledoux. Poincaré's inequalities and Talagrand's concentration phenomenon for the exponential distribution. *Probab. Theory Related Fields*, 107(3):383–400, 1997.
- [BLM13] S. Boucheron, G. Lugosi, and P. Massart. *Concentration inequalities*. Oxford University Press, Oxford, 2013. A nonasymptotic theory of independence, With a foreword by Michel Ledoux.
- [BTo6] S. G. Bobkov and P. Tetali. Modified logarithmic Sobolev inequalities in discrete settings. *J. Theoret. Probab.*, 19(2):289–336, 2006.
- [Buco1] A. Buchholz. Operator Khintchine inequality in non-commutative probability. *Math. Ann.*, 319(1):1–16, 2001.
- [Cafoo] L. A. Caffarelli. Monotonicity properties of optimal transportation and the FKG and related inequalities. *Comm. Math. Phys.*, 214(3):547–563, 2000.
- [CH16] H.-C. Cheng and M.-H. Hsieh. Characterizations of matrix and operator-valued Φ-entropies, and operator Efron-Stein inequalities. *Proc. A.*, 472(2187):20150563, 20, 2016.
- [CH19] H.-C. Cheng and M.-H. Hsieh. Matrix Poincaré, Φ-Sobolev inequalities, and quantum ensembles. *J. Math. Phys.*, 60(3):032201, 16, 2019.
- [Chao7] S. Chatterjee. Stein's method for concentration inequalities. Probab. Theory Related Fields, 138(1-2):305–321, 2007.
- [CHT17] H.-C. Cheng, M.-H. Hsieh, and M. Tomamichel. Exponential decay of matrix Φ-entropies on Markov semigroups with applications to dynamical evolutions of quantum ensembles. *J. Math. Phys.*, 58(9):092202, 24, 2017.
- [CT14] R. Y. Chen and J. A. Tropp. Subadditivity of matrix φ -entropy and concentration of random matrices. *Electron. J. Probab.*, 19:no. 27, 30, 2014.
- [Dav69] E. B. Davies. Quantum stochastic processes. Comm. Math. Phys., 15:277–304, 1969.
- [GKS20] A. Garg, T. Kathuria, and N. Srivastava. Scalar Poincaré implies matrix Poincaré. arXiv ma.PR 2006.09567, June
- [GLSS18] A. Garg, Y. T. Lee, Z. Song, and N. Srivastava. A matrix expander Chernoff bound. In STOC'18—Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing, pages 1102–1114. ACM, New York, 2018.
- [GM83] M. Gromov and V. D. Milman. A topological application of the isoperimetric inequality. *Amer. J. Math.*, 105(4):843–854, 1983.
- [Gro75] L. Gross. Hypercontractivity and logarithmic Sobolev inequalities for the Clifford Dirichlet form. *Duke Math. J.*, 42(3):383–396, 1975.
- [HS19] J. Hermon and J. Salez. Modified log-Sobolev inequalities for strong-Rayleigh measures, 2019.
- [HT20] D. Huang and J. A. Tropp. Nonlinear matrix concentration via semigroup methods. Manuscript, June 2020.
- [HZ15] F. Hansen and Z. Zhang. Characterisation of matrix entropies. Lett. Math. Phys., 105(10):1399-1411, 2015.
- [JZ15] M. Junge and Q. Zeng. Noncommutative martingale deviation and Poincaré type inequalities with applications. *Probab. Theory Related Fields*, 161(3-4):449–507, 2015.
- [KT13] M. J. Kastoryano and K. Temme. Quantum logarithmic Sobolev inequalities and rapid mixing. *J. Math. Phys.*, 54(5):052202, 30, 2013.
- [Ledo1] M. Ledoux. *The concentration of measure phenomenon*, volume 89 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2001.
- [Led97] M. Ledoux. On Talagrand's deviation inequalities for product measures. ESAIM Probab. Statist., 1:63–87, 1995/97.
- [Lin76] G. Lindblad. On the generators of quantum dynamical semigroups. Comm. Math. Phys., 48(2):119–130, 1976.
- [LP86] F. Lust-Piquard. Inégalités de Khintchine dans C_p (1 . C. R. Acad. Sci. Paris Sér. I Math., 303(7):289–292, 1986.
- [LT11] M. Ledoux and M. Talagrand. *Probability in Banach spaces*. Classics in Mathematics. Springer-Verlag, Berlin, 2011. Isoperimetry and processes, Reprint of the 1991 edition.
- [MJC⁺14] L. Mackey, M. I. Jordan, R. Y. Chen, B. Farrell, and J. A. Tropp. Matrix concentration inequalities via the method of exchangeable pairs. *Ann. Probab.*, 42(3):906–945, 2014.
- [MOZ98] A. W. Majewski, R. Olkiewicz, and B. Zegarlinski. Dissipative dynamics for quantum spin systems on a lattice. *J. Phys. A*, 31(8):2045–2056, 1998.
- [Mün19] F. Münch. Li–Yau inequality under cd(0, n) on graphs. arXiv ma.DG 1909.10242, Sep. 2019.

- [Olio9] R. I. Oliveira. Concentration of the adjacency matrix and of the Laplacian in random graphs with independent edges. 2009.
- [Oli10] R. I. Oliveira. Sums of random Hermitian matrices and an inequality by Rudelson. *Electron. Commun. Probab.*, 15:203–212, 2010.
- [OZ99] R. Olkiewicz and B. Zegarlinski. Hypercontractivity in noncommutative L_p spaces. *J. Funct. Anal.*, 161(1):246–285, 1999.
- [Pet94] D. Petz. A survey of certain trace inequalities. In *Functional analysis and operator theory (Warsaw, 1992)*, volume 30 of *Banach Center Publ.*, pages 287–298. Polish Acad. Sci. Inst. Math., Warsaw, 1994.
- [Pis86] G. Pisier. Probabilistic methods in the geometry of Banach spaces. In *Probability and analysis (Varenna, 1985*), volume 1206 of *Lecture Notes in Math.*, pages 167–241. Springer, Berlin, 1986.
- [PMT16] D. Paulin, L. Mackey, and J. A. Tropp. Efron–Stein inequalities for random matrices. *Ann. Probab.*, 44(5):3431–3473, 2016.
- [PP14] R. Pemantle and Y. Peres. Concentration of Lipschitz functionals of determinantal and other strong Rayleigh measures. *Combin. Probab. Comput.*, 23(1):140–160, 2014.
- [PV15] J. Pitrik and D. Virosztek. On the joint convexity of the Bregman divergence of matrices. *Lett. Math. Phys.*, 105(5):675–692, 2015.
- [PX97] G. Pisier and Q. Xu. Non-commutative martingale inequalities. Comm. Math. Phys., 189(3):667–698, 1997.
- [Rud99] M. Rudelson. Random vectors in the isotropic position. J. Funct. Anal., 164(1):60–72, 1999.
- [Str84] D. W. Stroock. An introduction to the theory of large deviations. Universitext. Springer-Verlag, New York, 1984.
- [Tro11] J. A. Tropp. Freedman's inequality for matrix martingales. *Electron. Commun. Probab.*, 16:262–270, 2011.
- [Tro12] J. A. Tropp. User-friendly tail bounds for sums of random matrices. Found. Comput. Math., 12(4):389-434, 2012.
- [Tro15] J. A. Tropp. An introduction to matrix concentration inequalities. *Foundations and Trends in Machine Learning*, 8(1-2):1–230, 2015.
- [Tro16] J. A. Tropp. The expected norm of a sum of independent random matrices: an elementary approach. In *High dimensional probability VII*, volume 71 of *Progr. Probab.*, pages 173–202. Springer, [Cham], 2016.
- [Var85] N. T. Varopoulos. Hardy-Littlewood theory for semigroups. J. Funct. Anal., 63(2):240–260, 1985.
- [VH16] R. Van Handel. Probability in High Dimension. APC 550 lecture notes, Princeton Univ., 2016.